

Babies, Toddlers and pre-school Vaccinations

6-in-one vaccine

Protects against: [diphtheria](#), [tetanus](#), [whooping cough](#), [polio](#), [Hib \(Haemophilus influenzae type b\)](#) and [hepatitis B](#).

Given at: 8, 12 and 16 weeks of age to all babies born on or after 1 August 2017

Pneumococcal or pneumo jab (PCV)

Protects against: some types of [pneumococcal infection](#)

Given at: 8 weeks, 16 weeks and one year of age

Rotavirus vaccine

Protects against: [rotavirus infection](#), a common cause of childhood diarrhoea and sickness

Given at: 8 and 12 weeks of age

Men B vaccine

Protects against: [meningitis](#) (caused by meningococcal type B bacteria)

Given at: 8 weeks, 16 weeks and one year of age

Hib/Men C vaccine

Protects against: [Haemophilus influenzae type b \(Hib\)](#) and [meningitis](#) caused by meningococcal group C bacteria

Given at: one year of age

MMR vaccine

Protects against: [measles](#), [mumps](#) and [rubella](#)

Given at: one year and at three years and four months of age

Children's flu vaccine

Protects against: [flu](#)

Given at: annually as a nasal spray in Sept/Oct for all children aged two to eight years on 31 August 2017

4-in-1 pre-school booster

Protects against: [diphtheria](#), [tetanus](#), [whooping cough](#) and [polio](#)

Given at: three years and four months of age

Babies, Toddlers and pre-school Vaccinations

These are the vaccines a child requires before they start Primary school

<u>Vaccination</u>	<u>Protection given</u>	<u>Age given at</u>
6-in-one vaccine	diphtheria , tetanus , whooping cough , polio , Hib (Haemophilus influenzae type b) and hepatitis B .	8, 12 and 16 weeks of age to all babies born on or after 1 August 2017
<u>Pneumococcal or pneumo jab (PCV)</u>	some types of pneumococcal infection	8 weeks, 16 weeks and one year of age
Rotavirus vaccine	rotavirus infection , a common cause of childhood diarrhoea and sickness	8 and 12 weeks of age
Men B vaccine	meningitis (caused by meningococcal type B bacteria)	8 weeks, 16 weeks and one year of age
Hib/Men C vaccine	Haemophilus influenzae type b (Hib) and meningitis caused by meningococcal group C bacteria	one year of age
MMR vaccine	measles , mumps and rubella	one year and at three years and four months of age
Children's flu vaccine	flu	Annually(Yearly) as a nasal spray in Sept/Oct for all children aged two to eight years on 31 August 2017
4-in-1 pre-school booster	diphtheria , tetanus , whooping cough and polio	three years and four months of age

Please note; Children's Flu vaccine is offered in school annually for children in Key stage 1 by the Immunisation Team from Central and North West London NHS Foundation Trust. You can read more about all these vaccinations on www.nhs.uk